

Preprint APCTP/98-22

KEK-TH-594

hep-ph/9810369

Decay Constants $f_{D_s^*}$ and f_{D_s} from $\bar{B}^0 \rightarrow D^+ l^- \bar{\nu}$ and $\bar{B}^0 \rightarrow D^+ D_s^{-(*)}$ Decays

Yong-Yeon Keum *

*Asia Pacific Center for Theoretical Physics**207-43 Cheongrangi Dong, Dongdaemun-Gu, Seoul 130-012, Korea**Theory Group, KEK, Tsukuba, Ibaraki 305-0801 Japan*

Abstract

We calculate the decay constant of D_s and D_s^* with $\bar{B}^0 \rightarrow D^+ \ell^- \nu$ and $\bar{B}^0 \rightarrow D^+ D_s^{-(*)}$ decays. In our analysis we assume the factorization ansatz and use two different form factor behaviours (constant and monopole-type) for $F_0(q^2)$. We also consider the QCD-penguin contributions in hadronic decays within the NDR renormalization scheme in a NLO calculation.

We estimate the decay constant of the D_s meson to be $219 \pm 46 \text{ MeV}$ for (pole/pole)-type form factor and $239 \pm 50 \text{ MeV}$ for (pole/constant)-type form factor. For D_s^* meson, we predict $f_{D_s^*} = 336 \pm 79 \text{ MeV}$, and $f_{D_s^*}/f_{D_s} = 1.41 \pm 0.41$ for (pole/constant)-type form factor.

PACS index : 12.15.-y, 13.20.-v, 13.25.Hw, 14.40.Nd, 14.65.Fy

Keywords : Factorization, Non-leptonic Decays, Decay Constant, Penguin Effects

*Monbushou Research Fellow

Email: keum@apctp.kaist.ac.kr, keum@theory.kek.jp

1. Introduction

Measuring purely leptonic decays of heavy mesons provides the cleanest way to determine the weak decay constants of heavy mesons, which connect the measured quantities, such as the $B\bar{B}$ mixing ratio, to CKM matrix elements V_{cb}, V_{ub} . However, at present it is not possible to determine f_B, f_{B_s}, f_{D_s} and $f_{D_s^*}$ experimentally from leptonic B and D_s decays. For instance, the decay rate for D_s^+ is given by [1]

$$\Gamma(D_s^+ \rightarrow \ell^+ \nu) = \frac{G_F^2}{8\pi} f_{D_s}^2 m_\ell^2 M_{D_s} \left(1 - \frac{m_\ell^2}{M_{D_s}^2}\right)^2 |V_{cs}|^2 \quad (1)$$

Because of helicity suppression, the electron mode $D_s^+ \rightarrow e^+ \nu$ has a very small rate. The relative widths are $10 : 1 : 2 \times 10^{-5}$ for $\tau^+ \nu, \mu^+ \nu$ and $e^+ \nu$ final states, respectively. Unfortunately the mode with the largest branching fraction, $\tau^+ \nu$, has at least two neutrinos in the final state and is difficult to detect experimentally. So, theoretical calculations for decay constant have to be used. The factorization ansatz for nonleptonic decay modes provides us a good approximate method to obtain nonperturbative quantities, i.e., form factors and decay constants which are hardly accessible in any other way [2, 3].

There are many ways that the quarks produced in a nonleptonic weak decay can arrange themselves into hadrons. The final state is linked to the initial state by complicated trees of gluon and quark interactions, pair production, and loops. These make the theoretical description of nonleptonic decays difficult. However, since the products of a B meson decay are quite energetic, it is possible that the complicated QCD interactions are less important and that the two quark pairs of the currents in the weak Hamiltonian, group individually into the final state mesons without further exchanges of gluons. The color-transparency argument suggests that a quark-antiquark pair remains a state of small size with a correspondingly small chromomagnetic moment until it is far from the other decay products.

Color transparency is the basis for the factorization hypothesis, in which amplitudes factorize into products of two current matrix elements. This ansatz is widely used in heavy quark physics, as it is almost the only way to treat hadronic decays.

In this paper we consider a way to determine weak decay constants f_{D_s} and $f_{D_s^*}$ under factorization ansatz. In section 2, we discuss how to extract the unknown parameter $|V_{cb} F_1^{BD}(0)|$

from the branching ratio of the semileptonic decay $\bar{B}^0 \rightarrow D^+ \ell^- \bar{\nu}$. In order to check the validity of the factorization assumption, we study the nonleptonic two body decays, for instance, $B \rightarrow D\rho, D\pi$ and $DK^{(*)}$ in section 3. In section 4, we discuss the determination of f_{D_s} and $f_{D_s^*}$ from $\bar{B}^0 \rightarrow D^+ D_s^{-(*)}$ decay modes. In our analysis, we consider the QCD-penguin effects which amount to about 11 % for $B \rightarrow DD_s$ and 5 % for $B \rightarrow DD_s^*$, which have been mentioned in Ref.[4].

2. Semileptonic Decay $\bar{B}^0 \rightarrow D^+ l^- \bar{\nu}$

From Lorentz invariance one finds the decomposition of the hadronic matrix element in terms of hadronic form factors:

$$\begin{aligned} \langle D^+(p_D) | J_\mu | \bar{B}^0(p_B) \rangle &= \left[(p_B + p_D)_\mu - \frac{m_B^2 - m_D^2}{q^2} q_\mu \right] F_1^{BD}(q^2) \\ &\quad + \frac{m_B^2 - m_D^2}{q^2} q_\mu F_0^{BD}(q^2), \end{aligned} \quad (2)$$

where $J_\mu = \bar{c}\gamma_\mu b$ and $q_\mu = (p_B - p_D)_\mu$. In the rest frame of the decay products, $F_1(q^2)$ and $F_0(q^2)$ correspond to 1^- and 0^+ exchanges, respectively. At $q^2 = 0$ we have the constraint

$$F_1^{BD}(0) = F_0^{BD}(0), \quad (3)$$

since the hadronic matrix element in (2) is nonsingular at this kinematic point.

The q^2 distribution in the semileptonic decay $\bar{B}^0 \rightarrow D^+ l^- \bar{\nu}$ is written in terms of the hadronic form factor $F_1^{BD}(q^2)$ as

$$\frac{d\Gamma(\bar{B}^0 \rightarrow D^+ l^- \bar{\nu})}{dq^2} = \frac{G_F^2}{24\pi^3} |V_{cb}|^2 [K(q^2)]^3 |F_1(q^2)|^2, \quad (4)$$

where the q^2 dependent momentum $K(q^2)$ is given by

$$K(q^2) = \frac{1}{2m_B} \left[(m_B^2 + m_D^2 - q^2)^2 - 4m_B^2 m_D^2 \right]^{1/2}. \quad (5)$$

In the zero lepton-mass limit, $0 \leq q^2 \leq (m_B - m_D)^2$.

For the q^2 dependence of the form factors, Wirbel *et al.* [8] assumed a simple pole formula for both $F_1(q^2)$ and $F_0(q^2)$ (we designate this scenario 'pole/pole'):

$$F_1(q^2) = F_1(0) / (1 - \frac{q^2}{m_{F_1}^2}), \quad F_0(q^2) = F_0(0) / (1 - \frac{q^2}{m_{F_0}^2}), \quad (6)$$

with the pole masses

$$m_{F_1} = 6.34 \text{ GeV}, \quad m_{F_0} = 6.80 \text{ GeV}. \quad (7)$$

Korner and Schuler [9] have also adopted the same q^2 dependence of $F_1(q^2)$ and $F_0(q^2)$ given by (6) and (7). On the other hand, the heavy quark effective theory (HQET) gives in the $m_{b,c} \rightarrow \infty$ limit the relation between $F_1(q^2)$ and $F_0(q^2)$ given by [11, 12]

$$F_0(q^2) = \left[1 - \frac{q^2}{(m_B + m_D)^2} \right] F_1(q^2). \quad (8)$$

The combination of (6) and (8) suggests that $F_0(q^2)$ is approximately constant when we keep the simple pole dependence for $F_1(q^2)$. Therefore, in this paper, as well as the above 'pole/pole' form factors, we will also consider the following ones (designated 'pole/const.')

$$F_1(q^2) = F_1(0) / (1 - \frac{q^2}{m_{F_1}^2}), \quad F_0(q^2) = F_0(0), \quad (9)$$

with

$$m_{F_1} = 6.34 \text{ GeV}. \quad (10)$$

By introducing the variable $x \equiv q^2/m_B^2$, which has the range of $0 \leq x \leq (1 - \frac{m_D}{m_B})^2$ in the zero lepton mass limit, (4) is written as

$$\begin{aligned} \frac{d\Gamma(\bar{B}^0 \rightarrow D^+ l^- \bar{\nu})}{dx} &= \frac{G_F^2 m_B^5}{192\pi^3} |V_{cb} F_1^{BD}(0)|^2 \frac{\lambda^3[1, \frac{m_D^2}{m_B^2}, x]}{\left(1 - \frac{m_D^2}{m_{F_1}^2} x\right)^2}, \\ \lambda[1, \frac{m_D^2}{m_B^2}, x] &= \left[\left(1 + \frac{m_D^2}{m_B^2} - x\right)^2 - 4 \frac{m_D^2}{m_B^2} \right]^{1/2}. \end{aligned} \quad (11)$$

Then the branching ratio $\mathcal{B}(\bar{B}^0 \rightarrow D^+ l^- \bar{\nu})$ is given by

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^+ l^- \bar{\nu}) &= \left(\frac{G_F m_B^2}{\sqrt{2}}\right)^2 \frac{m_B}{\Gamma_B} \frac{2}{192\pi^2} |V_{cb} F_1^{BD}(0)|^2 \times I \\ &= 2.221 \times 10^2 |V_{cb} F_1^{BD}(0)|^2 \times I, \end{aligned} \quad (12)$$

where the dimensionless integral I is given by

$$I = \int_0^{(1-\frac{m_D}{m_B})^2} dx \frac{\left[\left(1 + \frac{m_D^2}{m_B^2} - x\right)^2 - 4\frac{m_D^2}{m_B^2} \right]^{3/2}}{\left(1 - \frac{m_B^2}{m_{F_1}^2} x\right)^2} = 0.121 \quad (13)$$

In obtaining the numerical values in (12) and (13), we have used the following experimental results [33]: $m_D = m_{D^+} = 1.869$ GeV, $m_B = m_{B^0} = 5.279$ GeV, $\Gamma_B = \Gamma_{B^0} = 4.219 \times 10^{-13}$ GeV ($\tau_{B^0} = (1.56 \pm 0.06) \times 10^{-12}$ s), and $G_F = 1.166 \ 39(2) \times 10^{-5}$ GeV $^{-2}$. Since $\mathcal{B}(\bar{B}^0 \rightarrow D^+ l^- \bar{\nu}) = (1.78 \pm 0.20 \pm 0.24) \times 10^{-2}$ is obtained experimentally [13], the value of $|V_{cb} F_1^{BD}(0)|$ can be extracted from (12). Following this procedure, we obtain

$$|V_{cb} F_1^{BD}(0)| = (2.57 \pm 0.14 \pm 0.17) \times 10^{-2}. \quad (14)$$

In the calculations that follow, we will use $|V_{cb} F_1^{BD}(0)| = (2.57 \pm 0.22) \times 10^{-2}$ which combines the statistical and systematic errors in (14).

3. Test of Factorization with $\bar{B}^0 \rightarrow D^+ \rho^-$ and $\bar{B}^0 \rightarrow D^+ \pi^-$, and Prediction of Branching Ratio $\mathcal{B}(\bar{B}^0 \rightarrow D^+ K^{(*)})$

We start by recalling the relevant effective weak Hamiltonian:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* [C_1(\mu) \mathcal{O}_1 + C_2(\mu) \mathcal{O}_2] + \text{H.C.}, \quad (15)$$

$$\mathcal{O}_1 = (\bar{d}\Gamma^\rho u)(\bar{c}\Gamma_\rho b), \quad \mathcal{O}_2 = (\bar{c}\Gamma^\rho u)(\bar{d}\Gamma_\rho b), \quad (16)$$

where G_F is the Fermi coupling constant, V_{cb} and V_{ud} are corresponding Cabibbo-Kobayashi-Maskawa (CKM) matrix elements and $\Gamma_\rho = \gamma_\rho(1 - \gamma_5)$. The Wilson coefficients $C_1(\mu)$ and $C_2(\mu)$ incorporate the short-distance effects arising from the scaling of $\mathcal{H}_{\text{eff}}$ from $\mu = m_W$ to $\mu = O(m_b)$. By using the Fierz transformation under which $V - A$ currents remain $V - A$ currents, we get the following equivalent forms:

$$\begin{aligned} C_1 \mathcal{O}_1 + C_2 \mathcal{O}_2 &= (C_1 + \frac{1}{N_c} C_2) \mathcal{O}_1 + C_2 (\bar{d}\Gamma^\rho T^a u)(\bar{c}\Gamma_\rho T^a b) \\ &= (C_2 + \frac{1}{N_c} C_1) \mathcal{O}_2 + C_1 (\bar{c}\Gamma^\rho T^a u)(\bar{d}\Gamma_\rho T^a b), \end{aligned} \quad (17)$$

where $N_c = 3$ is the number of colors and T^a s are $SU(3)$ color generators. The second terms in (17) involve color-octet currents. In the factorization assumption, these terms are neglected and $\mathcal{H}_{\text{eff}}$ is rewritten in terms of “factorized hadron operators” [8, 10]:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* \left(a_1 [\bar{d} \Gamma^\rho u]_H [\bar{c} \Gamma_\rho b]_H + a_2 [\bar{c} \Gamma^\rho u]_H [\bar{d} \Gamma_\rho b]_H \right) + \text{H.C.}, \quad (18)$$

where the subscript H stands for *hadronic* implying that the Dirac bilinears inside the brackets be treated as interpolating fields for the mesons and no further Fierz-reordering need be done. The phenomenological parameters a_1 and a_2 are related to C_1 and C_2 by

$$a_1 = C_1 + \frac{1}{N_c} C_2, \quad a_2 = C_2 + \frac{1}{N_c} C_1. \quad (19)$$

From the analyses of A.J. Buras [14], the parameters a_1 and a_2 are determined in the Next-to-Leading-Order (NLO) calculation in the Naive Dimensional Regularization (NDR) scheme as

$$a_1 = 1.02 \pm 0.01, \quad a_2 = 0.20 \pm 0.05. \quad (20)$$

For the two body decay, in the rest frame of initial meson the differential decay rate is given by

$$d\Gamma = \frac{1}{32\pi^2} |\mathcal{M}|^2 \frac{|\mathbf{p}_1|}{M^2} d\Omega, \quad (21)$$

$$|\mathbf{p}_1| = \frac{[(M^2 - (m_1 + m_2)^2)(M^2 - (m_1 - m_2)^2)]^{1/2}}{2M}, \quad (22)$$

where M is the mass of initial meson, and m_1 (m_2) and $\mathbf{p}_1$ are the mass and momentum of one of final mesons. By using (2), (18) and $\langle 0 | \Gamma_\mu | \rho(q, \varepsilon) \rangle = \varepsilon_\mu(q) m_\rho f_\rho$, (21) gives the following formula for the branching ratio of $\bar{B}^0 \rightarrow D^+ \rho^-$:

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^+ \rho^-) &= \left(\frac{G_F m_B^2}{\sqrt{2}} \right)^2 |V_{ud}|^2 \frac{1}{16\pi} \frac{m_B}{\Gamma_B} a_1^2 \frac{f_\rho^2}{m_B^2} |V_{cb} F_1^{BD}(m_\rho^2)|^2 \\ &\times \left[\left(1 - \left(\frac{m_D + m_\rho}{m_B} \right)^2 \right) \left(1 - \left(\frac{m_D - m_\rho}{m_B} \right)^2 \right) \right]^{3/2} \\ &= 13.25 \times |V_{cb} F_1^{BD}(m_\rho^2)|^2. \end{aligned} \quad (23)$$

In obtaining the numerical values in (23), we have used $m_\rho = m_{\rho^+} = 766.9$ MeV, $f_\rho = f_{\rho^+} = 216$ MeV, $V_{ud} = 0.9751$ [33], and the parameters given below (13).

For a_1 we use the value given in (20). Then, by using the formula (23) with the values of $|V_{cb} F_0^{BD}(0)|^2$, ($F_0^{BD}(0) = F_1^{BD}(0)$) given in (14), we obtain the branching ratio $\mathcal{B}(\bar{B}^0 \rightarrow D^+ \rho^-)$ presented in Table 3.

For the process $\bar{B}^0 \rightarrow D^+ K^{*-}$, by using $\langle 0 | \Gamma_\mu | K^*(q, \varepsilon) \rangle = \varepsilon_\mu(q) m_{K^*} f_{K^*}$, we have

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^+ K^{*-}) &= \left(\frac{G_F m_B^2}{\sqrt{2}} \right)^2 |V_{us}|^2 \frac{1}{16\pi} \frac{m_B}{\Gamma_B} a_1^2 \frac{f_{K^*}^2}{m_B^2} |V_{cb} F_1^{BD}(m_{K^*}^2)|^2 \\ &\times \left[\left(1 - \left(\frac{m_D + m_{K^*}}{m_B} \right)^2 \right) \left(1 - \left(\frac{m_D - m_{K^*}}{m_B} \right)^2 \right) \right]^{3/2} \\ &= 0.67 \times |V_{cb} F_1^{BD}(m_{K^*}^2)|^2. \end{aligned} \quad (24)$$

where we have used $m_{K^*} = m_{K^{*-}} = 891.59$ MeV, $f_{K^*} = f_{K^{*-}} = 218$ MeV, and $V_{us} = 0.2215$ [33]. By using (23) with $|V_{cb} F_1^{BD}(0)|^2$ in (14), we obtain the branching ratio $\mathcal{B}(\bar{B}^0 \rightarrow D^+ K^{*-})$ presented in Table 1.

By using (2), (18) and $\langle 0 | \Gamma_\mu | \pi(q) \rangle = i q_\mu f_\pi$, (21) gives the following formula for the branching ratio for the process $\bar{B}^0 \rightarrow D^+ \pi^-$:

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^+ \pi^-) &= \left(\frac{G_F m_B^2}{\sqrt{2}} \right)^2 |V_{ud}|^2 \frac{1}{16\pi} \frac{m_B}{\Gamma_B} a_1^2 \frac{f_\pi^2}{m_B^2} |V_{cb} F_0^{BD}(m_\pi^2)|^2 \\ &\times \left(1 - \frac{m_D^2}{m_B^2} \right)^2 \left[\left(1 - \left(\frac{m_D + m_\pi}{m_B} \right)^2 \right) \left(1 - \left(\frac{m_D - m_\pi}{m_B} \right)^2 \right) \right]^{1/2} \\ &= 5.42 \times |V_{cb} F_0^{BD}(m_\pi^2)|^2. \end{aligned} \quad (25)$$

where we have used $m_\pi = m_{\pi^-} = 139.57$ MeV and $f_\pi = f_{\pi^-} = 131.74$ MeV [33]. By using the formula (25) with the values of $|V_{cb} F_0^{BD}(0)|^2$, ($F_0^{BD}(0) = F_1^{BD}(0)$) in (14), we obtain the branching ratio for $\bar{B}^0 \rightarrow D^+ \pi^-$ presented in Table 1.

For the process $\bar{B}^0 \rightarrow D^+ K^-$, by using $\langle 0 | \Gamma_\mu | K^-(q) \rangle = i q_\mu f_{K^-}$, we have

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^+ K^-) &= \left(\frac{G_F m_B^2}{\sqrt{2}} \right)^2 |V_{us}|^2 \frac{1}{16\pi} \frac{m_B}{\Gamma_B} a_1^2 \frac{f_K^2}{m_B^2} |V_{cb} F_0^{BD}(m_K^2)|^2 \\ &\times \left(1 - \frac{m_D^2}{m_B^2} \right)^2 \left[\left(1 - \left(\frac{m_D + m_K}{m_B} \right)^2 \right) \left(1 - \left(\frac{m_D - m_K}{m_B} \right)^2 \right) \right]^{1/2} \\ &= 0.41 \times |V_{cb} F_0^{BD}(m_K^2)|^2, \end{aligned} \quad (26)$$

where we have used $m_K = m_{K^-} = 493.68$ MeV, $f_K = f_{K^+} = 160.6$ MeV [33]. By using (23) with $|V_{cb} F_1^{BD}(0)|^2$ in (14), we obtain the branching ratio $\mathcal{B}(\bar{B}^0 \rightarrow D^+ K^-)$ presented in Table 1.

An inspection of Table 1 shows that the factorization method works well in $\bar{B}^0 \rightarrow D^+\pi^-, D^+\rho^-$ decays. We predict branching ratios :

$$\begin{aligned}\mathcal{B}(\bar{B}^0 \rightarrow D^+ K^-) &\simeq 2.7 \cdot 10^{-4} \\ \mathcal{B}(\bar{B}^0 \rightarrow D^+ K^{*-}) &\simeq 4.6 \cdot 10^{-4}\end{aligned}\quad (27)$$

which is certainly reachable in near future.

4. Determination of $f_{D_s^*}$ and f_{D_s} from $\bar{B}^0 \rightarrow D^+ D_s^{*-}$ and $\bar{B}^0 \rightarrow D^+ D_s^-$

(a) At Tree level

In this section we calculate the vector and pseudoscalar decay constants $f_{D_s^*}$ and f_{D_s} from the experimentally obtained branching ratios of the the exclusive decays $\bar{B}^0 \rightarrow D^+ D_s^{*-}$ and $\bar{B}^0 \rightarrow D^+ D_s^-$. By using (2), (18) and $\langle 0 | \Gamma_\mu | D_s^*(q, \varepsilon) \rangle = \varepsilon_\mu(q) m_{D_s^*} f_{D_s^*}$, (21) gives the following formula for the branching ratio of $\bar{B}^0 \rightarrow D^+ D_s^{*-}$:

$$\begin{aligned}\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^{*-}) &= \left(\frac{G_F m_B^2}{\sqrt{2}} \right)^2 |V_{cb}|^2 |V_{cs}|^2 \frac{1}{16\pi} \frac{m_B}{\Gamma_B} a_1^2 \frac{f_{D_s^*}^2}{m_B^2} |F_1^{BD}(m_{D_s^*}^2)|^2 \\ &\times \left[\left(1 - \left(\frac{m_D + m_{D_s^*}}{m_B} \right)^2 \right) \left(1 - \left(\frac{m_D - m_{D_s^*}}{m_B} \right)^2 \right) \right]^{3/2} \\ &= (1.32 \cdot 10^2 \text{ GeV}^2) \cdot f_{D_s^*}^2 \cdot |V_{cb} F_1^{BD}(m_{D_s^*}^2)|^2,\end{aligned}\quad (28)$$

where we have used $m_{D_s^*} = m_{D_s^{*-}} = 2112.4 \text{ MeV}$ $V_{cs} = 0.9743$ [33]. Then, from (28) we get

$$f_{D_s^*} = (0.87 \times 10^{-1} \text{ GeV}) \cdot \frac{\sqrt{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^{*-})}}{|V_{cb} F_1^{BD}(m_{D_s^*}^2)|}.\quad (29)$$

For the process $\bar{B}^0 \rightarrow D^+ D_s^-$, by using $\langle 0 | \Gamma_\mu | D_s(q) \rangle = i q_\mu f_{D_s}$, we have

$$\begin{aligned}\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-) &= \left(\frac{G_F m_B^2}{\sqrt{2}} \right)^2 |V_{cb}|^2 |V_{cs}|^2 \frac{1}{16\pi} \frac{m_B}{\Gamma_B} a_1^2 \frac{f_{D_s}^2}{m_B^2} |F_0^{BD}(m_{D_s}^2)|^2 \\ &\times \left(1 - \frac{m_D^2}{m_B^2} \right)^2 \left[\left(1 - \left(\frac{m_D + m_{D_s}}{m_B} \right)^2 \right) \left(1 - \left(\frac{m_D - m_{D_s}}{m_B} \right)^2 \right) \right]^{1/2} \\ &= (2.45 \cdot 10^2 \text{ GeV}^2) \cdot f_{D_s}^2 \cdot |V_{cb} F_0^{BD}(m_{D_s}^2)|^2,\end{aligned}\quad (30)$$

where we used $m_{D_s} = m_{D_s^-} = 1968.5$ MeV [33]. From (30) we get

$$f_{D_s} = (0.64 \times 10^{-1} \text{ GeV}) \cdot \frac{\sqrt{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-)}}{|V_{cb} F_0^{BD}(m_{D_s}^2)|}. \quad (31)$$

Browder *et al.* [15] present the following experimental results for the branching ratios:

$$\begin{aligned} \mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^{*-}) &= (1.14 \pm 0.42 \pm 0.28) \times 10^{-2} = (1.14 \pm 0.50) \times 10^{-2}, \\ \mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-) &= (0.74 \pm 0.22 \pm 0.18) \times 10^{-2} = (0.74 \pm 0.28) \times 10^{-2}, \end{aligned} \quad (32)$$

where we have combined the statistical and systematic errors. By using (32) and the values of $|V_{cb} F_1^{BD}(m_{D_s^*}^2)|$ and $|V_{cb} F_0^{BD}(m_{D_s}^2)|$ given in Table 2, from (31) and (29) we obtain the following results:

$$\begin{aligned} f_{D_s^*} &= 322 \pm 76 \text{ MeV}, & f_{D_s} &= 196 \pm 41 \text{ MeV} & \text{for (pole/pole),} \\ f_{D_s^*} &= 322 \pm 76 \text{ MeV}, & f_{D_s} &= 214 \pm 45 \text{ MeV} & \text{for (pole/const.).} \end{aligned} \quad (33)$$

The ratio of the vector and pseudoscalar decay constants $f_{D_s^*}/f_{D_s}$ is given by

$$\frac{f_{D_s^*}}{f_{D_s}} = 1.36 \times \frac{|V_{cb} F_0^{BD}(m_{D_s}^2)|}{|V_{cb} F_1^{BD}(m_{D_s^*}^2)|} \times \left[\frac{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^{*-})}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-)} \right]^{1/2}, \quad (34)$$

which gives

$$\begin{aligned} \frac{f_{D_s^*}}{f_{D_s}} &= 1.64 \pm 0.48 & \text{for (pole/pole),} \\ \frac{f_{D_s^*}}{f_{D_s}} &= 1.51 \pm 0.44 & \text{for (pole/const.).} \end{aligned} \quad (35)$$

(b) Including Penguin contribution

The effective Hamiltonian for $\Delta B = 1$ transitions is given by

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} [V_{ub} V_{uq}^* (C_1 O_1^u + C_2 O_2^u) + V_{cb} V_{cq}^* (C_1 O_1^c + C_2 O_2^c) - V_{tb} V_{tq}^* \sum_{i=3}^6 C_i O_i], \quad (36)$$

where $q = d, s$ and C_i are the Wilson coefficients evaluated at the renormalization scale μ , and the current-current operators $O_1^{u,c}$ and $O_2^{u,c}$ are

$$\begin{aligned} O_1^u &= (\bar{u}_\alpha b_\alpha)_{V-A} (\bar{q}_\beta u_\beta)_{V-A} & O_1^c &= (\bar{c}_\alpha b_\alpha)_{V-A} (\bar{q}_\beta c_\beta)_{V-A} \\ O_2^u &= (\bar{u}_\beta b_\alpha)_{V-A} (\bar{q}_\alpha u_\beta)_{V-A} & O_2^c &= (\bar{c}_\beta b_\alpha)_{V-A} (\bar{q}_\alpha c_\beta)_{V-A}, \end{aligned} \quad (37)$$

and the QCD penguin operators $O_3 - O_6$ are

$$\begin{aligned} O_3 &= (\bar{q}_\alpha b_\alpha)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\beta)_{V-A} & O_4 &= (\bar{q}_\beta b_\alpha)_{V-A} \sum_{q'} (\bar{q}'_\alpha q'_\beta)_{V-A} \\ O_5 &= (\bar{q}_\alpha b_\alpha)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\beta)_{V+A} & O_6 &= (\bar{q}_\beta b_\alpha)_{V-A} \sum_{q'} (\bar{q}'_\alpha q'_\beta)_{V+A}. \end{aligned} \quad (38)$$

In (36) we have neglected the effects of the electroweak penguin operators and the dipole operators, since their contributions are not important for the calculations of this paper.

When we take $m_t = 174$ GeV, $m_b = 5.0$ GeV, $\alpha_s(M_z) = 0.118$ and $\alpha_{\text{em}}(M_z) = 1/128$, the numerical values of the renormalization scheme independent Wilson coefficients $\bar{C}_i$ at $\mu = m_b$ are given by [16]

$$\begin{aligned} \bar{C}_1 &= -0.3125, & \bar{C}_2 &= 1.1502, \\ \bar{C}_3 &= 0.0174, & \bar{C}_4 &= -0.0373, & \bar{C}_5 &= 0.0104, & \bar{C}_6 &= -0.0459. \end{aligned} \quad (39)$$

The effective Hamiltonian in (36) for the decays $\bar{B}^0 \rightarrow D^+ D_s^{-(*)}$ can be rewritten as

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} [V_{cb} V_{cs}^* (C_1^{\text{eff}} O_1^c + C_2^{\text{eff}} O_2^c) - V_{tb} V_{ts}^* \sum_{i=3}^6 C_i^{\text{eff}} O_i], \quad (40)$$

where C_i^{eff} are given by [17]

$$\begin{aligned} C_1^{\text{eff}} &= \bar{C}_1, & C_2^{\text{eff}} &= \bar{C}_2, & C_3^{\text{eff}} &= \bar{C}_3 - P_s/N_c, & C_4^{\text{eff}} &= \bar{C}_4 + P_s, \\ C_5^{\text{eff}} &= \bar{C}_5 - P_s/N_c, & C_6^{\text{eff}} &= \bar{C}_6 + P_s \end{aligned} \quad (41)$$

with

$$\begin{aligned} P_s &= \frac{\alpha_s}{8\pi} \left[\frac{10}{9} - G(m_q, q^2, \mu) \right] \bar{C}_2(\mu), \\ G(m_q, q^2, \mu) &= -4 \int_0^1 x(1-x) \ln \left(\frac{m_q^2 - x(1-x)q^2}{\mu^2} \right) dx, \end{aligned} \quad (42)$$

where q denotes the momentum of the virtual gluons appearing in the QCD time-like matrix elements, and N_c is the number of colors. Assuming $q^2 = m_b^2/2$, we obtain the analytic formular for $G(m_q, q^2, \mu)$:

$$G(m_q, \frac{m_b^2}{2}, \mu) = -\frac{2}{3} \ln \left(\frac{y}{8} \right) + \frac{10}{9} + \frac{2}{3} y + \frac{(2+y)\sqrt{1-y}}{3} \left[\ln \left| \frac{1-\sqrt{1-y}}{1+\sqrt{1-y}} \right| + i\pi \right] \quad (43)$$

with $y = 8m_q^2/m_b^2$. By inserting the values for $m_q = m_c = 1.45 \text{ GeV}$, and $N_c = 3$ into (41) and (43), we get the values $C_i^{eff}(i = 1 \sim 6)$ which is shown in Table 2.

The decay amplitude $\mathcal{A}_{T+P}(\bar{B}^0 \rightarrow D^+ D_s^-) = \langle D^+ D_s^- | \mathcal{H}_{\text{eff}} | \bar{B}^0 \rangle$ is given as follows:

$$\begin{aligned} \mathcal{A}_{T+P}(\bar{B}^0 \rightarrow D^+ D_s^-) &= \frac{G_F}{\sqrt{2}} [V_{cb} V_{cs}^* a_2 - V_{tb} V_{ts}^* (a_4 + 2a_6 \frac{m_{D_s}^2}{(m_b - m_c)(m_s + m_c)})] \mathcal{M}_a \\ &\simeq \frac{G_F}{\sqrt{2}} V_{cb} V_{cs}^* a_2 \left[1 + \frac{a_4}{a_2} + 2 \frac{a_6}{a_2} \frac{m_{D_s}^2}{(m_b - m_c)(m_s + m_c)} \right] \mathcal{M}_a \end{aligned} \quad (44)$$

where

$$\mathcal{M}_a = \langle D_s^- | \bar{s} \gamma^\mu \gamma_5 c | 0 \rangle \langle D^+ | \bar{c} \gamma_\mu b | \bar{B}^0 \rangle = -i f_{D_s} (m_B^2 - m_D^2) F_0^{BD}(m_{D_s}^2) \quad (45)$$

On the other hand, we have

$$\begin{aligned} \mathcal{A}_{T+P}(\bar{B}^0 \rightarrow D^+ D_s^{*-}) &= \frac{G_F}{\sqrt{2}} [V_{cb} V_{cs}^* a_2 - V_{tb} V_{ts}^* a_4] \mathcal{M}_b \\ &\simeq \frac{G_F}{\sqrt{2}} V_{cb} V_{cs}^* a_2 \left(1 + \frac{a_4}{a_2} \right) \mathcal{M}_b \end{aligned} \quad (46)$$

where

$$\mathcal{M}_b = \langle D_s^* | \bar{s} \gamma^\mu \gamma_5 c | 0 \rangle \langle D^+ | \bar{c} \gamma_\mu b | \bar{B}^0 \rangle = m_{D_s^*} f_{D_s^*} [\epsilon(q) \cdot (P_B + P_D)] F_1^{BD}(m_{D_s^*}^2) \quad (47)$$

We can estimate the penguin contributions for each process from (44) and (46) as follows :

$$\text{For } \bar{B}^0 \rightarrow D^+ D_s^-; \quad \left| \frac{\mathcal{A}_P}{\mathcal{A}_T} \right| = \left| \frac{a_4}{a_2} + 2 \frac{a_6}{a_2} \frac{m_{D_s}^2}{(m_b - m_c)(m_c + m_s)} \right| = 10.9\% \quad (48)$$

$$\text{For } \bar{B}^0 \rightarrow D^+ D_s^{*-}; \quad \left| \frac{\mathcal{A}_P}{\mathcal{A}_T} \right| = \left| \frac{a_4}{a_2} \right| = 4.5\% \quad (49)$$

Here we have used the values for $m_c = 1.45 \text{ GeV}$, and $m_s = 170 \text{ MeV}$. The penguin contributions for $B \rightarrow DD_s$ is two times larger than that in $B \rightarrow DD_s^*$, and is a sizable effect. This results agrees well with that of Ref.[4]. Hence penguin contributions can affect the value of the decay constants f_{D_s} and $f_{D_s^*}$.

From (31), (29), (44) and (46) we obtain the following results including penguin contributions :

$$\begin{aligned} f_{D_s^*} &= 336 \pm 79 \text{ MeV}, & f_{D_s} &= 219 \pm 46 \text{ MeV} & \text{for (pole/pole),} \\ f_{D_s^*} &= 336 \pm 79 \text{ MeV}, & f_{D_s} &= 239 \pm 50 \text{ MeV} & \text{for (pole/const.).} \end{aligned} \quad (50)$$

From (34), (44) and (46) the ratio of the vector and pseudoscalar decay constants $f_{D_s^*}/f_{D_s}$ is given by

$$\frac{f_{D_s^*}}{f_{D_s}} = 1.36 \cdot \frac{|V_{cb} F_0^{BD}(m_{D_s}^2)|}{|V_{cb} F_1^{BD}(m_{D_s}^2)|} \cdot \left[\frac{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^{*-})}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-)} \right]^{1/2} \cdot \left(\frac{0.8951}{0.9582} \right) \quad (51)$$

which gives

$$\begin{aligned} \frac{f_{D_s^*}}{f_{D_s}} &= 1.53 \pm 0.45 \quad \text{for (pole/pole),} \\ \frac{f_{D_s^*}}{f_{D_s}} &= 1.41 \pm 0.41 \quad \text{for (pole/const.).} \end{aligned} \quad (52)$$

The decay constant depends the q^2 behaviour of the form factor $F_0(q^2)$. However the amount of change is less than 10% as shown in (50). This shows that the decay constant is not strongly dependent on the behaviour of the form factor. As shown in Table 3, when we include the QCD-penguin contributions, the value of the decay constant $f_{D_s^*}$ is increased by 4%; however, for f_{D_s} it is increased by up to 12%, and the ratio $f_{D_s^*}/f_{D_s}$ is decreased by 6%. Table 3 shows that our result for f_{D_s} agrees well with others' predictions and experimental data, except that of Hwang and Kim [18, 19]. For the ratio $f_{D_s^*}/f_{D_s}$, our results have a value greater than 1. On the contrary, Browder et al. [15] and Hwang and Kim 2 [19] have a value less than 1, which is quite strange, because their results show that the decay constant of the vector meson is smaller than that of the pseudoscalar meson with the same quark contents. In fact, the decay constant of ρ meson is 1.5 times greater than that of π meson. By measuring the ratio $f_{D_s^*}/f_{D_s}$, we can check directly their analysis and methods.

5. Conclusion

Within the factorization approximation, we have calculated the weak decay constants f_{D_s} and $f_{D_s^*}$ from the semileptonic decay, $\bar{B}^0 \rightarrow D^+ \ell^- \nu$ and the hadronic decay modes, $\bar{B}^0 \rightarrow D^+ \bar{D}_s^{(*)}$. We have also considered the effect of two different q^2 -dependences of the form factor $F_0^{BD}(q^2)$. The value of f_{D_s} is altered by less than 10% for different form factors. In our analysis, we also considered the QCD-penguin contributions in hadronic two body decays within the NDR renormalization scheme in a next-to-leading order calculation. The penguin effects for $B \rightarrow DD_s$ decay are quite sizable. We obtain $f_{D_s} = 219 \pm 46 \text{ MeV}$ for the monopole type of F_0^{BD} , $f_{D_s} = 239 \pm 50 \text{ MeV}$ for the constant F_0^{BD} . We obtain $f_{D_s^*} = 336 \pm 79 \text{ MeV}$ for the

D_s^* meson. These predicted values will be improved vastly when the large accumulated data samples is available at Belle and BaBar experiment in near future.

Acknowledgements

The author wishes to thank D. S. Hwang for helpful discussions. He is grateful to A. N. Kamal for reading this manuscript carefully. Y.-Y. K. would like to thank M. Kobayashi for his hospitality and encouragement. This work is supported in part by the Basic Science Research Institute Program, Ministry of Education, Project No. BSRI-97-2414, and in part by the Grant-in Aid for Scientific from the Ministry of Education, Science and Culture, Japan.

References

- [1] J. L. Rosner, in **Particles and Fields 3**, Proceeding of the 1988 Banff Summer Inst., Banff, Alberta, Canada, ed. by A.N. Kamal and F.C. Khanna, World Scientific. Singapore, 1989, 395.
- [2] D. Bortoletto and S. Stone, Phys. Rev. Lett. 65 (1990) 2951.
- [3] M. Neubert, V. Rieckert, B. Stech and Q. P. Xu, in **Heavy Flavours** ed. by A. J. Buras and M. Lindner, World Scientific. Singapore, 1992
- [4] M. Gourdin, A. N. Kamal, Y.Y. Keum and X. Y. Pham, Phys. Lett.B 333 (1994) 507.
- [5] CLEO Collaboration, J.P. Alexander et al., Phys. Rev. Lett. 77 (1996) 5000.
- [6] CLEO Collaboration, D.M. Asner et al., Phys. Rev. D 53 (1996) 1039.
- [7] A. Ali, Z. Phys. C 1 (1979) 25.

- [8] M. Wirbel, B. Stech and M. Bauer, Z. Phys. C 29 (1985) 637; M. Bauer and M. Wirbel, Z. Phys. C 42 (1989) 671.
- [9] J.G. Korner and G.A. Schuler, Z. Phys. C 38 (1988) 511; *ibid.* (erratum) C 41 (1989) 690.
- [10] Y.-Y. Keum, Phys. Lett. B 348 (1995) 247.
- [11] N. Isgur and M.B. Wise, Phys. Lett. B 232 (1989) 113; *ibid.* B 237 (1990) 527.
- [12] M. Neubert and V. Rieckert, Nucl. Phys. B 382 (1992) 97.
- [13] S. Stone, hep-ph/9610305, in Proceedings of NATO Advanced Study Institute on Techniques and Concepts of High Energy Physics, Virgin Islands, July 1996.
- [14] A. J. Buras, Nucl. Phys. B **434**, 606 (1995).
- [15] T.E. Browder, K. Honscheid and D. Pedrini, hep-ph/9606354, UH-515-848-96, OHSTPY-HEP-E-96-006, To appear in *Annual Review of Nuclear and Particle Science*, Vol. 46.
- [16] N.G. Deshpande and X.-G. He, Phys. Rev. Lett. **74**, 26 (1995).
- [17] R. Fleischer, Z. Phys. **C58**, 483 (1993); **C62**, 81 (1994).
- [18] D.S. Hwang and G.-H. Kim, Phys. Rev. D **55**, 6944 (1997).
- [19] D.S. Hwang and G.-H. Kim, in Proceedings of the Third International Workshop on Particle Physics Phenomenology, Taipei, Taiwan, Nov. 1996 (World Scientific, Singapore).
- [20] S. Capstick and S. Godfrey, Phys. Rev. D **41**, 2856 (1990).
- [21] C.A. Dominguez, in Proceedings of the Third Workshop on Tau-Charm Factory, Marbella, Spain, June, 1993.
- [22] R.M. Boxter *et al.*, UKQCD Collaboration, Phys. Rev. D **49**, 1594 (1994).
- [23] C.W. Bernard, J.N. Labrenz and A. Soni, Phys. Rev. D **49**, 2536 (1994).
- [24] C. Bernard *et al.*, MILC Collaboration, hep-ph/9709328 (1997).

- [25] S. Aoki *et al.*, WA75 Collaboration, Prog. Theor. Phys. **89**, 131 (1993).
- [26] D. Acosta *et al.*, CLEO Collaboration, Phys. Rev. D **49**, 5690 (1994).
- [27] D. Gibaut *et al.*, CLEO Collaboration, CLEO CONF 95-22, EPS0184 (1995).
- [28] J.Z. Bai *et al.*, BES Collaboration, Phys. Rev. Lett. **74**, 4599 (1995).
- [29] K. Kodama *et al.*, E653 Collaboration, Phys. Lett. **B382**, 299 (1996).
- [30] B. Grinstein, M.B. Wise and N. Isgur, Phys. Rev. Lett. 56 (1986) 298; N. Isgur, D. Scora, B. Grinstein and M.B. Wise, Phys. Rev. D 39 (1989) 799 (1989).
- [31] M. Suzuki, Phys. Rev. D 37 (1988) 239.
- [32] UKQCD Collaboration, D.R. Burford *et al.*, Nucl. Phys. B 447 (1995) 425.
- [33] Review of Particle Physics, R.M. Barnett *et al.*, Phys. Rev. D **54**, 1 (1996).
- [34] M. Chadha *et al.*, CLEO Collaboration, Phys. Rev. D **58**, 32002 (1998).

	$\mathcal{B}(\bar{B}^0 \rightarrow D^+ \rho^-)$ $\times 10^3$	$\mathcal{B}(\bar{B}^0 \rightarrow D^+ K^{*-})$ $\times 10^4$	$\mathcal{B}(\bar{B}^0 \rightarrow D^+ \pi^-)$ $\times 10^3$	$\mathcal{B}(\bar{B}^0 \rightarrow D^+ K^-)$ $\times 10^4$
(pole/pole)	9.01 ± 1.54	4.62 ± 0.79	3.58 ± 0.61	2.74 ± 0.47
(pole/const.)	9.01 ± 1.54	4.62 ± 0.79	3.57 ± 0.61	2.71 ± 0.46
Experiments	$8.4 \pm 1.6 \pm 0.7$	—	$3.1 \pm 0.4 \pm 0.2$	—

Table 1: The obtained values of the branching ratios and existing experimental values.

Coefficients	Real Part	Imaginary Part
C_1^{eff}	- 0.313	0.0
C_2^{eff}	1.150	0.0
C_3^{eff}	$2.20 \cdot 10^{-2}$	$5.13 \cdot 10^{-3}$
C_4^{eff}	$-5.11 \cdot 10^{-2}$	$-1.54 \cdot 10^{-2}$
C_5^{eff}	$1.50 \cdot 10^{-2}$	$5.13 \cdot 10^{-3}$
C_6^{eff}	$-5.97 \cdot 10^{-2}$	$-1.54 \cdot 10^{-2}$
a_1	0.071	0.0
a_2	1.046	0.0
a_3	$4.97 \cdot 10^{-3}$	0.0
a_4	$-4.38 \cdot 10^{-2}$	$-1.37 \cdot 10^{-2}$
a_5	$-4.90 \cdot 10^{-3}$	0.0
a_6	$-5.47 \cdot 10^{-2}$	$-1.37 \cdot 10^{-2}$

Table 2: The values of the effective wilson coefficient C_i^{eff} and a_i with the $\mu = m_b = 5.0 \text{ GeV}$, $m_c = 1.45 \text{ GeV}$ in NDR scheme at NLO calculation. The coefficients a_{2i} and a_{2i-1} are defined by $a_{2i-1} = C_{2i-1}^{eff} + C_{2i}^{eff}/N_c$, $a_{2i} = C_{2i}^{eff} + C_{2i-1}^{eff}/N_c$ and we have taken $N_c = 3$.

	$f_{D_s^*}$ (MeV)	f_{D_s} (MeV)	$f_{D_s^*}/f_{D_s}$
(pole/pole) at Tree level	322 ± 76	196 ± 41	1.64 ± 0.48
(pole/const.) at Tree level	322 ± 76	214 ± 45	1.51 ± 0.44
(pole/pole) with Penguin	336 ± 79	219 ± 46	1.53 ± 0.45
(pole/const.) with Penguin	336 ± 79	239 ± 50	1.41 ± 0.41
Browder <i>et al.</i> [15]	243 ± 70	277 ± 77	0.88 ± 0.35
Hwang and Kim 1 [18]	362 ± 15	309 ± 15	1.17 ± 0.02
Hwang and Kim 2 [19]	336 ± 13	399 ± 24	0.84 ± 0.03
Capstick and Godfrey [20]		290 ± 20	
Dominguez [21]		222 ± 48	
UKQCD [22]		212^{+4+46}_{-3-7}	
BLS [23]		$230 \pm 7 \pm 35$	
MILC [24]		$199 \pm 8^{+40+10}_{-11-0}$	
WA75 [25]		$238 \pm 47 \pm 21 \pm 48$	
CLEO 1 [26]		$282 \pm 30 \pm 43 \pm 34$	
CLEO 2 [27]		$280 \pm 19 \pm 28 \pm 34$	
BES [28]		$430^{+150}_{-130} \pm 40$	
E653 [29]		$190 \pm 34 \pm 20 \pm 26$	

Table 3: The obtained values of $f_{D_s^*}$ (MeV) and f_{D_s} (MeV), and their ratio $f_{D_s^*}/f_{D_s}$, and the results from other theoretical calculations and existing experimental results. Here we referred the corrected f_{D_s} values [34] for the experimental data [25] - [29].